



Contents lists available at ScienceDirect

Journal of Oral and Maxillofacial Surgery, Medicine, and Pathology

journal homepage: www.elsevier.com/locate/jomsmp



Original Research

Evaluation of environmental change in the mouth with the use of spray-type oral moisturizer containing γ -PGA



Shoichi Yamamoto^{a,*}, Hiroaki Yoshida^a, Tadashi Ohkubo^b, Hirohumi Sawai^b, Shosuke Morita^a

^a First Department of Oral and Maxillofacial Surgery, Osaka Dental University, 1-5-17, Otemae, Chuo-ku, Osaka 540-0008, Japan

^b Department of Internal Medicine, Osaka Dental University, 1-5-17, Otemae, Chuo-ku, Osaka 540-0008, Japan

ARTICLE INFO

Article history:

Received 23 March 2016

Received in revised form 23 May 2016

Accepted 8 June 2016

Available online 27 July 2016

Keywords:

Xerostomia

Spray-type oral moisturizer

γ -PGA

Dry mouth

Stress marker

ABSTRACT

Objective: This study was to evaluate the effect of new spray-type oral moisturizer containing poly- γ -glutamic acid (γ -PGA). γ -PGA can be classified as a pseudopolyamino acid, which contains only repeated glutamate units. The hygroscopic and moisturizing effects of γ -PGA were compared with HA. Cross-linked γ -PGA could absorb thousands of times more water than its own weight.

Methods: 102 volunteers were randomly allocated to experimental and control groups. The experimental group (50 subjects) used this new moisturizer, the control group I (20 subjects) used sterile distilled water at normal temperature, and the control group II (32 subjects) used nothing. In experimental group, the effect of moisturizer was investigated by measuring amylase activity (salivary amylase monitor[®]), oral moisture (Mucus[®]), stimulated salivary flow rates at predetermine times (Saxon test).

Results: In the amylase activity and the oral moisture, there was not a significant difference in all groups. In the stimulated salivary flow rates, there was a significant difference. When we compared the experimental group with the control group I, an increase was admitted at after 30 min ($p < 0.05$). When we compared the experimental group with the control group II, an increase was admitted at after 10 ($p < 0.01$), 20 ($p < 0.05$), 30 min ($p < 0.05$).

Conclusions: New moisturizer was able to improve the environment in the mouth. Oral application of γ -PGA has a possibility to be a therapeutic strategy for treating Xerostomia.

© 2016 Asian AOMS, ASOMP, JSOP, JSOMS, JSOM, and JAMI. Published by Elsevier Ltd. All rights reserved. *

1. Introduction

Saliva is an important body fluid that plays a vital role in protecting the oral mucosa and teeth, as well as aiding oral function and the digestion of food. An absence of saliva results in a number of oral changes and related behaviors that can influence a patient's quality of life [1–5].

Xerostomia induces bacteria propagation in the oral environment making it vulnerable to pneumonia caused by aspiration of saliva. Treatment includes reduction of stress and drug treatment such as herbal medicine and salivary gland hormones. Systemic therapies include taste stimulus treatment [6], massage of

salivary glands [7], gum treatment [8], and using a moisturizing agent [9–11].

In Japan, Aquabalance[®] medical mouthspray was commercialized as the first spray-type oral moisturizer containing γ -PGA by LION Dental Products Company in 2013.

Poly- γ -glutamic acid (γ -PGA) can be classified as a pseudopolyamino acid, which contains only repeated glutamate units. However, γ -PGA differs from proteins, both structurally and functionally. The glutamate units in γ -PGA are linked between the α -amino and γ -carboxylic acid functional groups, whereas the peptide linkages in proteins are formed between the amino and γ -carboxylic acid groups. *Bacillus subtilis* (natto) was isolated from the traditional Japanese food, natto, Fermented soybean health food containing abundant γ -PGA, which is popular in Japan. Forming a highly mucous colony, the strain can grow on a high-saline medium containing L-glutamic acid. *Bacillus subtilis* (natto) is a plasmid-free γ -PGA producer [12].

The purpose of this study is analysis of the oral environmental changes with this spray.

* Asian AOMS: Asian Association of Oral and Maxillofacial Surgeons; ASOMP: Asian Society of Oral and Maxillofacial Pathology; JSOP: Japanese Society of Oral Pathology; JSOMS: Japanese Society of Oral and Maxillofacial Surgeons; JSOM: Japanese Society of Oral Medicine; JAMI: Japanese Academy of Maxillofacial Implants.

* Corresponding author. Tel.: +81 6 6910 1076; fax: +81 6 6910 1028.

E-mail address: yamamo-s@cc.osaka-dent.ac.jp (S. Yamamoto).

2. Materials and methods

2.1. Subjects

The subjects were 102 healthy Japanese students ranging from 20 to 28 years old (46 males and 56 females) at Osaka Dental University. All subjects gave written informed consent to participation in this study after receiving explanation of its purpose. The experimental group was 50 subjects (27 males and 23 females) who had Aquabalance medical mouthspray® sprayed on the back of the tongue. The control group I was 20 subjects (5 males and 15 females) who had sprayed sterile distilled water at normal temperature on the back of the tongue. The control group II was 32 subjects (14 males and 18 females) who did not receive spray application.

2.2. Oral moisturizer

Aquabalance medical mouthspray® (AB) used in this study is a spray-type oral moisturizer containing l-menthol, γ -poly glutamic acid (*Bacillus natto* gum) [13], glycerin, propylene glycol, polyoxyethylene hydrogenated castor oil (60E.O.), xylitol, buffering agent, flavor, and cetylpyridinium chloride.

2.3. Study design

This study was undertaken with the approval of the Institutional Review Board (IRB) of the Osaka Dental University Hospital (IRB number: 110837).

2.3.1. Examination 1: Measurement of amylase activity by salivary amylase monitor

Amylase was measured with a salivary amylase monitor (Nipro Co., Japan) and test strip (Nipro Co., Japan) [14,15]. We used this measurement method because it was easy and less invasive, since the test strip was placed under the tongue for only 30 s and also because it only required about 1 min from saliva sampling until obtaining the result.

The salivary amylase monitor builds on a precursor colorimetric assay platform wherein the color intensity of the enzymatic reaction product is photometrically measured to determine the concentration of salivary α -amylase. The biosensor system comprises of a disposable test strip and a hand-held reader with digital display. The test strip integrates a collector pad at its tip and a reagent paper infused with Gal-G2-CNP, a chromogenic substrate for salivary α -amylase. When placed under the tongue, the collector pad quickly saturates with fixed micro-liter quantities of saliva. The strip is inserted into the reader and the lever rises to activate the reader and transfer the collected saliva onto the reagent paper. After a 10 s interval, an audible signal indicates completion of saliva transfer and the strip is retracted. The salivary amylase activity in the transferred saliva begins metabolizing the Gal-G2-CNP substrate to yield the colored product CNP according to the reaction. Enzymatic activity is allowed to proceed for 10 s and the reflectance of the reaction product then measured photometrically at 430 nm. The color measured is proportional to the concentration of the salivary amylase activity; the greater the intensity of the color observed, the greater the concentration of salivary amylase activity. The biosensor's micro-processing unit calculates the salivary amylase activity level and displays it as a digital readout along with a date and time stamp [16,17].

Table 1
Measuring of amylase activity by salivary amylase monitor.

Group of subjects	Experimental	Control I		Control II	
		Mean	SD	Mean	SD
Baseline	21.68			37.42	
After 10 min	20.17			38.79	NS
After 20 min	24.67			36.56	NS
After 30 min	18.73			35.94	NS
				25.21	NS
				24.74	NS

2.3.2. Examination 2: Measurement of oral moisture by Mucus® [18]

Oral moisture was measured at the lingual mucosa (10 mm from the apex linguae on the surface of the tongue) and at the right and left buccal mucosa (10 mm from the angle of the mouth).

2.3.3. Examination 3: Measurement of stimulated salivary flow rates by Saxon test [19]

Saliva production was measured by weighing a folded sterile gauze pad before and after chewing. The low-normal value is 2 g/2 min.

The subjects were randomly allocated to one of the three groups:

Experimental group: 50 persons who used AB.

Control group I: 20 persons who did nothing.

Control group II: 32 persons who used sterile distilled water.

We performed all examinations in baseline (before application them) after 10, 20, and 30 min for experimental and control group.

We carried out this study between noon and 3 pm and measured after more than 1 h of eating and/or drinking. During the experiment, eating, drinking, and conversation were forbidden. The subjects were told to relax in a sitting position.

2.4. Statistical analysis

We compared the experimental group with the control group I and II with *T*-test. We compared baseline after 10, 20, and 30 min on each group.

3. Results

3.1. Examination 1

We measured amylase activity by salivary amylase monitor.

There was not a significant difference between Experimental group and Control group I for 30 min.

There was not a significant difference between Experimental group and Control group II for 30 min (Table 1).

3.2. Examination 2

We measured oral moisture by Mucus.

There was not a significant difference between Experimental group and Control group I for 30 min.

Table 2
Measuring of the oral moisture by Mucus®.

Group of subjects	Experimental	Control 1		Control 2	
		Mean	SD	Mean	SD
Baseline	1.95			2.2	
After 10 min	2.69			1.65	NS
After 20 min	2.3			1.7	NS
After 30 min	1.99			1.85	NS
				1.2	NS
				1.85	NS

Table 3

Measuring of the stimulated salivary flow rates by Saxon test.

Group of subjects	Experimental	Control 1		Control 2	
		Mean	SD	Mean	SD
Baseline	1.43	1.63		2.03	
After 10 min	1.68	1.69	NS	2.34	$p < 0.01$
After 20 min	1.37	1.81	NS	2.36	$p < 0.05$
After 30 min	1.27	1.67	$p < 0.05$	2.34	$p < 0.05$

There was not a significant difference between Experimental group and Control group II for 30 min (Table 2).

3.3. Examination 3

We measured stimulated salivary flow rates by Saxon test.

There was a significant difference between Experimental group and Control group I after 30 min.

There was a significant difference between Experimental group and Control group II for 30 min (Table 3).

When we compared the experimental group with the control group I, an increase was admitted at after 30 min ($p < 0.05$).

When we compared the experimental group with the control group II, an increase was admitted at after 10 ($p < 0.01$), 20 ($p < 0.05$), 30 min ($p < 0.05$).

4. Discussion

Saliva production and secretion is vital for maintenance of oral health and oral function. The complications of Xerostomia arise due to the effect of the oral secretions on microbial colonization of the oral cavity, demineralization and diminished remineralization of teeth, reduction in denture retention, mucosal dehydration and atrophy, reduced clearance rates of substances from the mouth (including microbes and food products), reduction in antibody and enzyme effects in the oral environment, and reduced lubrication of the oral tissues [20]. Common complications include rampant dental caries, candidiasis, mucosal atrophy and burning, difficulty in denture retention and use, compromised speech and swallowing, and reduced or altered taste sensation.

There are a lot of methods to measure mental and physical health. Especially, it is well known that glucocorticoid, catecholamine, amylase, cortisol and chromogranin A are stress markers [21,22]. Glucocorticoid and catecholamine require blood sample, and could cause stress. On the other hand, amylase and chromogranin A, which need saliva for inspection, react sensitively to an acute stress load. However, cortisol needs more than 10 min to react. There is a small quantity of chromogranin A in the saliva and there is individual variation. So we selected amylase as a stress marker in this study. The result of Examination 1 did not detect significant difference. We considered that there were no changes of the oral environment from the result of Examination 1 because this study was made for healthy persons.

Saliva measuring sheet and Mucus® are available to check oral moisture. This measures percent weight of water in the oral mucosal epithelium with the capacitance from the dielectric constant. Water content and the dielectric constant are positively correlated, and as the dielectric constant of water is markedly higher than that of other substances, the percentage of water in the substance can be determined by measuring the dielectric constant of the substance. The usefulness and ease of use of this oral hygrometer (within 2 s) have been reported. Following the manufacturer's definitions, oral moistness over 29% was considered normal, 25–29% was considered dry, and <25% was considered severely dry. Each measurement was performed in triplicate, and the median values were used in the subsequent analysis [23,24].

Within the normal range of the secreting saliva, the saliva measuring sheet does not react sometimes because the saliva viscosity and the food residue affect result of the saliva measuring sheet. So, we used Mucus® in this study.

The result of Examination 2 did not detect significant difference. We considered that there was no change of the moisture in an orally healthy person.

Saxon test and gum test were popular methods for measurement of saliva secretion quantity for outpatients. However, gum test requires more than 10 min. And so, Saxon test was conducted in this study. In Examination 3, there was a significant difference of stimulated whole salivary flow rate after application of AB.

In this study, there is a possibility that γ -PGA has a saliva secretion as these results.

Jae-Chul et al. reported that ultra-high molecular weight γ -PGA promotes wound healing in vivo [25].

Furthermore, Tae-Young et al. reported that oral administration of γ -PGA could be a therapeutic strategy for treating atopic dermatitis via the modulation of Th2-biased immune responses in an Nc/Nga mouse model [26].

Moon-Hee et al. reported that γ -PGA is well known as both a hydrophilic and a biodegradable macropolymer, which has high viscosity. γ -PGA is utilized in the manufacture of thickeners and cosmetics. As a moisturizer in cosmetics, γ -PGA can substitute for hyaluronic acid (HA), which is a carbohydrate or more specifically a mucopolysaccharide occurring naturally throughout the human body. HA can be several thousands of sugar (carbohydrate) molecules long. It is one of the most hydrophilic molecules in nature and can be described as "nature's moisturizer," but it is very expensive. In late 2002, the hygroscopic and moisturizing effects of γ -PGA were compared with HA. Cross-linked γ -PGA could absorb thousands of times more water than its own weight [12].

In conclusion, we detected that the environment in the mouth was improved with the use of AB spray-type oral moisturizer containing γ -PGA. It might play an important role in the maintenance of a healthy oral environment.

Ethical approval

The human rights and individual information of the subjects were protected and managed according to the ethical guidelines of Institutional Review Board of the Osaka Dental University Hospital (Approval No. from the Institutional Review Board: 110837).

Conflict of interest

All authors declare that there are no conflicts of interests.

References

- [1] Fox PC, van der Ven PF, Sonies BC, Weiffenbach JM, Baum BJ. Xerostomia: evaluation of a symptom with increasing significance. *J Am Dent Assoc* 1985;110:519–25.
- [2] FDI Working Group 10, CORE. Saliva: its role in health and disease. *Int Dent J* 1992;42:291–304.
- [3] Locker D. Subjective reports of oral dryness in an older adult population. *Community Dent Oral Epidemiol* 1993;21:165–8.
- [4] Guggenheim J, Moore PA. Xerostomia: etiology, recognition and treatment. *J Am Dent Assoc* 2003;134:61–9.
- [5] Atkinson JC, Grisius M, Massey W. Salivary hypofunction and xerostomia: diagnosis and treatment. *Dent Clin North Am* 2005;49:309–26.
- [6] Lee JM, Garon E, Wong DT. Salivary diagnostics. *Orthod Craniofac Res* 2009;12:206–11.
- [7] Agha-Hosseini F, Mirzaii-Dizgah I, Moghaddam PP, Akrad ZT. Stimulated whole salivary flow rate and composition in menopausal women with oral dryness feeling. *Oral Dis* 2007;13:320–3.
- [8] Rejnfeldt I, Andersson P, Renvert S. Oral health status in individuals with dementia living in special facilities. *Int J Dent Hyg* 2006;4:67–71.

- [9] Epstein JB, Emerton S, Le ND, Stevenson-Moore P. A double-blind crossover trial of Oral Balance gel and Biotene1 toothpaste versus placebo in patients with xerostomia following radiation therapy. *Oral Oncol* 1999;35:132–7.
- [10] Makita Y, Hirose T, Sekine M, Fujiwara S, Wang PL. Relationship between sensory evaluation and rheological properties of oral moisturizers. *J Hard Tissue Biol* 2013;22:319–24.
- [11] Shahdad SA, Taylor C, Barclay SC, Steen IN, Preshaw PM. A double-blind, crossover study of Biotene Oral Balance and Bioxtra systems as salivary substitutes in patients with post-radiotherapy xerostomia. *Eur J Cancer Care* 2005;14:319–26.
- [12] Moon-hee S, Chung P, Chul-joong K, Haryoung P, Kenji S, Makoto A. Natural and edible biopolymer poly- γ -glutamic acid: synthesis, production, and applications. *Chem Rec* 2005;5:352–66.
- [13] Yu-jin K, Kyung-jin M, Jin-hwa H, Jae-Kwan L. Efficacy of poly-gamma-glutamic acid in women with high-risk human papillomavirus-positive vaginal intraepithelial neoplasia: an observational pilot study. *J Microbiol Biotechnol* 2015;25:1163–9.
- [14] Masaki Y, Mitsuo D, Junichi W, Shin O, Noriyasu T, Tomoyuki H. Hand-held monitor of sympathetic nervous system using salivary amylase activity and its validation by driver fatigue assessment. *Biosens Bioelectron* 2006;21:1007–14.
- [15] Kazuki H, Ryuji K, Kyota O, Yumiko T. Influence of flow experience during daily life on health-related quality of life and salivary amylase activity in Japanese college students. *Japan Soc Occup Med Traumatol* 2011;59:13–8.
- [16] Vivek S, Corwin Z, Theodore FR, David E, Masaki Y. Developmental validation of a point-of-care, salivary α -amylase biosensor. *Psychoneuroendocrinology* 2011;36:193–9.
- [17] Masaki Y, Takahiro K, Masashi K, Noriyasu T, Yasufumi M, Hiroshi Y. Performance evaluation of salivary amylase activity monitor. *Biosens Bioelectron* 2004;20:491–7.
- [18] Yamada H, Nakagawa Y, Nomura Y, Yamamoto K, Suzuki M, Watanabe NY. Preliminary results of moisture checker for mucus in diagnosing dry mouth. *Oral Dis* 2005;11:405–7.
- [19] Suh KI, Lee JY, Chung JW, Kim YK, Kho HS. Relationship between salivary flowrate and clinical symptoms and behaviours in patients with dry mouth. *J Oral Rehabil* 2007;34:739–44.
- [20] Epstein JB, Scully C. The role of saliva in oral health and causes and effects of xerostomia. *J Can Dent Assoc* 1992;58:217–21.
- [21] Rivera Go'mez B, Herna'ndez Vallejo G, Arriba de la Fuente L, López Cantor M, Díaz M, López Pintor RM. The relationship between the levels of salivary cortisol and the presence of xerostomia in menopausal women. A preliminary study. *Med Oral Patol Oral Cir Bucal* 2006;11:E407–12.
- [22] Noriyasu T, Masaki Y, Toshiaki A, Kenji E, Kenji U, Yasuo N. Effect of psychological stress on the salivary cortisol and amylase levels in healthy young adults. *Arch Oral Biol* 2004;49:963–8.
- [23] Takahashi F, Takahashi M, Toya S, Koji T, Morita O. Clinical usefulness of an oral moisture checking device (Mucus). *Prosthodont Res Pract* 2006;5:214–8.
- [24] Murakami M, Nishi Y, Kamashita Y, Nagaoka E. Relationship between medical treatment and oral dryness diagnosed by oral moisture-checking device in patients with maxillofacial prostheses. *J Prosthodont Res* 2009;53:67–71.
- [25] Jae-Chul C, Hiroshi U, chul-hoon L, Moon-hee S. Promotion effects of ultra-high molecular weight poly- γ -glutamic acid on wound healing. *J Microbiol Biotechnol* 2015;25:941–5.
- [26] Tae-Young L, Doo-jin K, Ji-na W, Il-han L, Moon-hee S, Haryoung P. Oral administration of poly- γ -glutamate ameliorates atopic dermatitis in Nc/Nga mice by suppressing Th2-biased immune response and production of IL-17A. *J Invest Dermatol* 2014;134:704–11.